


Listing of the claims


This listing of claims provided below will replace all prior versions.

1. (currently amended) A method of ~~achieving an immunomodulatory effect, achieving an antineoplastic effect, or~~ inhibiting hyperproliferative cell growth in a patient in need thereof, comprising administering to said patient an effective amount of a compound of formula I ~~formulae I to XVII~~ or a pharmaceutically acceptable salt thereof



wherein,

- B is a phenyl ring,
- D is a phenyl ring ~~or a 5-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2 or 3 heteroatoms selected from O, S, and N,~~
- A is, in each case independently of each other, a 5-membered saturated or partially or fully unsaturated heterocyclic ring containing 1 ~~[[,]]~~ or 2 ~~or 3~~ heteroatoms selected from O, S, and N,
- Q is a bond or an alkylene or alkenylene group containing 1-5 carbon atoms, which is optionally substituted with =O, and in which optionally a carbon atom is replaced with an N atom,
- Z₁ is, in each case independently, -NH₂, =O, =NH, ~~or =N-phenyl, -phenyl, or alkyl~~ containing 1 to 5 carbon atoms,
- Z₂ is, in each case independently, -OH, halogen, alkyl containing 1-5 carbon atoms, or CO₂H ~~which is optionally substituted with halogen, and/or substituted with =O and/or -OH, and in which one C atom is optionally replaced with an O atom,~~
- Z₃ is, in each case independently, alkyl containing 1-5 carbon atoms, and
- n is, in each case independently, 0, 1, 2, or 3~~[[,]]~~



wherein,

G_1 , G_2 , and G_3 are, in each case independently, C, O, S, or N;

D is a phenyl ring or a 5- or 6-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2 or 3 heteroatoms selected from O, S, and N;


Q is a straight chain or branched alkylene or alkenylene group containing 1-5 carbon atoms, which is optionally substituted with =O and/or -OH, and in which optionally 1 or 2 carbon atoms, independently of each other, are replaced with an N or S atom;

Z_1 is, in each case independently, =O, halogen, or an alkyl group containing 1-5 carbon atoms, which is optionally substituted with =O and/or -OH;

Z_2 is, in each case independently, =O, or an alkyl group containing 1-5 carbon atoms, which is optionally substituted with =O and/or -OH;

n is 0, 1, or 2; and

m is 0, or 1;



wherein,

G_1 , G_2 , G_3 , G_4 , G_5 and G_6 are, in each case independently, C, O, S, or N, such that four or five of G_1 , G_2 , G_3 , G_4 , G_5 and G_6 are C atoms and the remaining G_1 , G_2 , G_3 , G_4 , G_5 and G_6 are O, S, or N;

Q is a bond or a straight chain or branched alkylene or alkenylene group containing 1-10

carbon atoms which is optionally substituted with =O in one or two places, and in which optionally 1 or 2 carbon atoms, independently of each other, are replaced with an N or O atom, and in which optionally a carbon atom is replaced with a 6-membered heterocyclic group containing 1 or 2 nitrogen atoms when the alkylene or alkenylene group is a straight chain group;


Z_1 is, in each case independently, OH, halogen, or an alkyl group containing 1-5 carbon atoms;

Z_2 is, in each case independently, =O, halogen, or an alkyl group containing 1-10 carbon atoms which is optionally substituted with =O in one or two places and/or OH, and in which optionally 1 or 2 carbon atoms, independently of each other, are replaced with an N and/or S atom;

Z_3 is, in each case independently, OH, halogen, NO_2 , an alkyl group containing 1-5 carbon atoms, in which optionally a carbon atom is replaced with an O atom, and which is optionally substituted with =O in one or two places, or is O-phenyl, wherein the phenyl group in the O-phenyl is optionally substituted with an NO_2 group;

n is 0, 1, or 2, and

m is 0, 1, 2, or 3;



wherein,

B is a phenyl ring;

D_1 is a phenylene ring or a 6-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2, or 3 heteroatoms selected from O, S, and N;


D_2 and D_2' are, each independently of each other, absent or a phenyl or phenylene ring or a 5- or 6-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2 or 3 heteroatoms selected from O, S, and N;

~~Q₁ is a bond or a branched or straight chain alkylene or alkenylene group containing 1-10 carbon atoms, which is optionally substituted with 1 to 5 =O and/or OH groups, in which optionally 1, 2, or 3 carbon atoms are, in each case independently, replaced with an N, O or S atom, wherein S is optionally substituted with 1 or 2 =O groups,~~

~~Q₂ and Q₂' are, each independently of each other, a bond or a branched or straight chain alkylene group containing 1-5 carbon atoms, which is optionally substituted with an =O group, in which optionally a carbon atom is replaced with an N, S, or O atom, wherein Q₂ is absent when D₂ is absent and Q₂' is absent when D₂' is absent,~~

~~Z is, in each case independently, =O, =S, OH, NH₂, NO₂, C≡N, SO₃H, is halogen, or a straight chain or branched alkyl or alkenyl group containing 1 to 10, which is optionally substituted with 1 to 3 =O and/or OH groups, and in which optionally a carbon atom is replaced with an N, O or S atom, or is a cyclic alkyl group containing 3 carbon atoms,~~

~~n is, in each case independently, 0, 1, 2, 3, 4 or 5;~~




wherein,

~~Z is, in each case independently, NO₂, an alkyl containing 1 to 5 carbon atoms, in which optionally a carbon atom is replaced with an O atom, and which is optionally substituted with an =O group,~~

~~A is a straight chain alkylene group containing 1 to 5 carbon atoms, and~~

~~n is 1, 2 or 3;~~



wherein,

B is a phenyl ring,

D is absent, or is a phenyl ring or a 6-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2, or 3 heteroatoms selected from O, S, and N;

A is a 5-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms selected from O, S, and N;


Q₁ and Q₂ are, in each case independently of each other, a bond or a straight chain or branched alkylene group containing 1-5 carbon atoms, in which optionally a carbon atom is replaced with an O, N or S atom, and in which optionally 1 or 2 C groups are replaced with C= or =C groups, and which is optionally substituted with an =O group; wherein Q₂ is absent when D is absent;

Z₁ is, in each case independently, NO₂, OH, halogen, or an alkyl group containing 1-5 carbon atoms, which is optionally substituted with up to 3 =O and/or OH groups, and in which optionally a carbon atom is replaced with an N, S or O atom;

Z₂ is, in each case independently, NH₂, OH, =NH, =O, =S, phenyl, halogen, or an alkyl group containing 1-5 carbon atoms, which is optionally substituted with up to 3 =O and/or OH groups, and in which optionally a carbon atom is replaced with an S atom;

Z₃ is, in each case independently, =O, OH, NO₂, NH₂, halogen, or an alkyl group containing 1-5 carbon atoms, which is optionally substituted with up to 3 =O and/or OH groups, and in which optionally a carbon atom is replaced with an O atom, and

n is, in each case independently, 0, 1, 2 or 3;



wherein,

G_1 to G_7 are, in each case independently, C, O, S, or N, wherein at least 3 of G_1 to G_7 are C atoms,


Z_1 is, in each case independently, absent, or =O, =NH or an alkyl group containing 1 to 5 carbon atoms,

Z_2 is, in each case independently, a straight chain or branched alkyl group containing 1 to 5 carbon atoms, in which optionally a carbon atom is replaced with an O atom, and which is optionally substituted with 1 or 2 =O and/or OH groups,

Q is, in each case independently, a bond or an alkylene group containing 1 to 5 carbon atoms, which is optionally substituted with =O, in which optionally 1, 2, or 3 carbon atoms are, in each case independently, replaced with an N or S atom, wherein S is optionally substituted with 1 or 2 =O groups, and

n is 0, 1 or 2, and

m is 1 or 2;



wherein,

A is a 5 membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2 or 3 heteroatoms selected from O, S, and N, or is a C_{10} aromatic bi-cyclic ring containing 1, 2 or 3 heteroatoms selected from O, S, and N,

~~D is absent or is a fully or partially saturated or unsaturated cyclic ring containing 6 or 7 carbon atoms;~~


~~Q₁ and Q₂ are, each independently of each other, a bond or a straight chain or branched alkylene group containing 1-10 carbon atoms, which is optionally substituted with an =O group, and in which optionally 1, 2 or 3 carbon atoms, independently of each other, are replaced with an N or O atom, and wherein optionally 1-3 carbon atoms are replaced with a C= and/or =C, and/or when the alkylene group is straight chain with a phenyl group, wherein Q₂ is absent when D is absent;~~

~~Z₁ is, in each case independently, absent or an alkyl group containing 1 to 5 carbon atoms, in which optionally a carbon atom is replaced with an O group, and which is optionally substituted with one or two =O or OH groups;~~

~~Z₂ is, in each case independently, =O or an alkyl group containing 1-5 carbon atoms, which is optionally substituted with =O in one or two places and/or OH;~~

~~Z₃ is halogen, or an alkyl group containing 1 to 5 carbon atoms, which is optionally halogenated, and~~

~~n is 1 or 2;~~




~~wherein;~~

~~A is a 5 or 6 membered saturated or partially or fully unsaturated heterocyclic ring containing 2 or 3 heteroatoms selected from S and N;~~

~~Z is, in each case independently, a straight chain or branched alkyl group containing 3-5 carbon atoms, which is substituted with =O and/or OH groups, and in which a carbon atom is replaced with an S atom, and~~

~~n is 1, 2, or 3;~~



wherein,

G_1 to G_{12} are, each independently of each other, C, N, S or O,


Z is, in each case independently, an alkyl containing 1 to 5 carbon atoms, which is optionally substituted with 1 to 2 =O and/or -OH groups,

Q is a bond or an alkylene group containing 1 to 5 carbon atoms,

m 0, 1, 2 or 3,

n 0, 1, 2 or 3, such that

$m+n \geq 1$;



wherein,


Z_1 is, in each case independently, halogen, NO_2 or -OH,

Z_2 is, in each case independently, an alkyl group containing 1-5 carbon atoms, which is optionally substituted with an =O and/or -OH group, and in which optionally a carbon atom is replaced with an S atom,

n is 0, 1, 2, or 3,

m is 0, 1, 2, or 3, and


$n+m$ is 3 or more;



wherein,

~~Z is, in each case independently, C≡N, halogen, or an alkyl group containing 1-5 carbon atoms, which is optionally substituted with halogen, and/or is substituted with one or more =O and/or -OH groups, and in which optionally a carbon atom is replaced with an S atom, and~~

~~n is 2, 3, 4 or 5;~~



wherein,

~~D₁ is a 5- or 6- membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2, or 3 heteroatoms selected from O, S, and N;~~

~~D₂ is a 6- membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2, or 3 heteroatoms selected from O, S, and N, or is optionally a phenylene group when D₃ is present;~~

~~D₃ is absent or a 5- or 6- membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2, or 3 heteroatoms selected from O, S, and N;~~

~~Q₁ is -O-, or a straight chain alkylene group containing 1-5 carbon atoms, in which optionally a carbon atom is replaced with an N, O or S atom, and which is optionally substituted with an =O atom;~~


~~Q₂ is absent when D₃ is absent or is a bond or an -O- group;~~

~~Z₁ is, in each case independently, =O or halogen;~~

~~Z₂ is, in each case independently, =O, C≡N, COOH, NO₂ or halogen;~~

~~Z₃ is, in each case independently, halogen, and is absent when D₃ is absent, and~~

n is, in each case independently, 0, 1, 2, or 3;



wherein,


B is a phenylene group,

Q is a straight chain alkylene group containing 1-10 carbon atoms, in which optionally up to three carbon atoms are replaced with an N, O or S atom, and which is optionally substituted with 1 or 2 =O groups,

Z is, in each case independently, halogen, or an alkyl group containing 1-5 carbon atoms, in which optionally a carbon atom is replaced with an O atom,

n is 0 or 1, and

m is 1 or 2;




wherein,

D is, a 6-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2, or 3 heteroatoms selected from O, S, and N,

Z is =O

n is 1, or 2;




wherein,

Q is, each independently, C or N, wherein,

Z_1 is a phenyl group, or 2 of Z_1 together form with the Q atoms to which they are bound a 6-membered aromatic ring containing only C atoms,

Z_2 is halogen, preferably Cl, and

n is 1, or 2;




wherein,

D is, a carbocyclic group containing 8 to 10 carbon atoms, and


R is OH or an alkyl group containing 1 to 5 carbon atoms, in which optionally a carbon atom is replaced with an N or O atom or with a phenyl group, and which is optionally substituted with 1 to 2 =O and/or OH groups.

2. (currently amended) A method of inhibiting ~~modulating~~ the binding of a p56^{lck} molecule via an SH2 domain thereof to a corresponding cellular binding protein, or inhibiting ~~modulating~~ the activity of a p56^{lck} molecule via an SH2 domain thereof, comprising administering a compound of formula I ~~formula I to XVII~~ or a pharmaceutically acceptable salt thereof



wherein,

- B is a phenyl ring,
- D is a phenyl ring ~~or a 5-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2 or 3 heteroatoms selected from O, S, and N,~~
- A is, in each case independently of each other, a 5-membered saturated or partially or fully unsaturated heterocyclic ring containing 1 ~~[[,]]~~ or 2 ~~or 3~~ heteroatoms selected from O, S, and N,
- Q is a bond or an alkylene or alkenylene group containing 1-5 carbon atoms, which is optionally substituted with =O, and in which optionally a carbon atom is replaced with an N atom,
- Z₁ is, in each case independently, ~~NH₂, =O, =NH, or =N~~ ~~phenyl, phenyl, or alkyl~~ containing 1 to 5 carbon atoms,
- Z₂ is, in each case independently, ~~OH, halogen, alkyl containing 1-5 carbon atoms, or CO₂H which is optionally substituted with halogen, and/or substituted with =O and/or OH, and in which one C atom is optionally replaced with an O atom,~~
- Z₃ is, in each case independently, alkyl containing 1-5 carbon atoms, and
- n is, in each case independently, 0, 1, 2, or 3 ~~[[,]]~~



wherein,

G_1 , G_2 , and G_3 are, in each case independently, C, O, S, or N;

D is a phenyl ring or a 5- or 6-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2 or 3 heteroatoms selected from O, S, and N;


Q is a straight chain or branched alkylene or alkenylene group containing 1-5 carbon atoms, which is optionally substituted with =O and/or -OH, and in which optionally 1 or 2 carbon atoms, independently of each other, are replaced with an N or S atom;

Z_1 is, in each case independently, =O, halogen, or an alkyl group containing 1-5 carbon atoms, which is optionally substituted with =O and/or -OH;

Z_2 is, in each case independently, =O, or an alkyl group containing 1-5 carbon atoms, which is optionally substituted with =O and/or -OH;

n is 0, 1, or 2, and

m is 0, or 1;



wherein,

G_1 , G_2 , G_3 , G_4 , G_5 and G_6 are, in each case independently, C, O, S, or N, such that four or five of G_1 , G_2 , G_3 , G_4 , G_5 and G_6 are C atoms and the remaining G_1 , G_2 , G_3 , G_4 , G_5 and G_6 are O, S, or N;

Q is a bond or a straight chain or branched alkylene or alkenylene group containing 1-10

carbon atoms which is optionally substituted with =O in one or two places, and in which optionally 1 or 2 carbon atoms, independently of each other, are replaced with an N or O atom, and in which optionally a carbon atom is replaced with a 6-membered heterocyclic group containing 1 or 2 nitrogen atoms when the alkylene or alkenylene group is a straight chain group;


Z_1 is, in each case independently, OH, halogen, or an alkyl group containing 1-5 carbon atoms;

Z_2 is, in each case independently, =O, halogen, or an alkyl group containing 1-10 carbon atoms which is optionally substituted with =O in one or two places and/or OH, and in which optionally 1 or 2 carbon atoms, independently of each other, are replaced with an N and/or S atom;

Z_3 is, in each case independently, OH, halogen, NO_2 , an alkyl group containing 1-5 carbon atoms, in which optionally a carbon atom is replaced with an O atom, and which is optionally substituted with =O in one or two places, or is O-phenyl, wherein the phenyl group in the O-phenyl is optionally substituted with an NO_2 group;

n is 0, 1, or 2, and

m is 0, 1, 2, or 3;



wherein,

B is a phenyl ring;

D_1 is a phenylene ring or a 6-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2, or 3 heteroatoms selected from O, S, and N;


D_2 and D_2' are, each independently of each other, absent or a phenyl or phenylene ring or a 5- or 6-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2 or 3 heteroatoms selected from O, S, and N;

Q_1 — is a bond or a branched or straight chain alkylene or alkenylene group containing 1-10 carbon atoms, which is optionally substituted with 1 to 5 =O and/or OH groups, in which optionally 1, 2, or 3 carbon atoms are, in each case independently, replaced with an N, O or S atom, wherein S is optionally substituted with 1 or 2 =O groups,

Q_2 and Q_2' — are, each independently of each other, a bond or a branched or straight chain alkylene group containing 1-5 carbon atoms, which is optionally substituted with an =O group, in which optionally a carbon atom is replaced with an N, S, or O atom, wherein Q_2 is absent when D_2 is absent and Q_2' is absent when D_2' is absent,

Z — is, in each case independently, =O, =S, OH, NH_2 , NO_2 , $C\equiv N$, SO_3H , is halogen, or a straight chain or branched alkyl or alkenyl group containing 1 to 10, which is optionally substituted with 1 to 3 =O and/or OH groups, and in which optionally a carbon atom is replaced with an N, O or S atom, or is a cyclic alkyl group containing 3 carbon atoms,

n — is, in each case independently, 0, 1, 2, 3, 4 or 5;




wherein,

Z — is, in each case independently, NO_2 , an alkyl containing 1 to 5 carbon atoms, in which optionally a carbon atom is replaced with an O atom, and which is optionally substituted with an =O group,

A — is a straight chain alkylene group containing 1 to 5 carbon atoms, and

n — is 1, 2 or 3;



wherein,

B is a phenyl ring,

D is absent, or is a phenyl ring or a 6-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2, or 3 heteroatoms selected from O, S, and N,

A is a 5-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms selected from O, S, and N,


Q₁ and Q₂ are, in each case independently of each other, a bond or a straight chain or branched alkylene group containing 1-5 carbon atoms, in which optionally a carbon atom is replaced with an O, N or S atom, and in which optionally 1 or 2 C groups are replaced with C= or =C groups, and which is optionally substituted with an =O group, wherein Q₂ is absent when D is absent,

Z₁ is, in each case independently, NO₂, OH, halogen, or an alkyl group containing 1-5 carbon atoms, which is optionally substituted with up to 3 =O and/or OH groups, and in which optionally a carbon atom is replaced with an N, S or O atom,

Z₂ is, in each case independently, NH₂, OH, =NH, =O, =S, phenyl, halogen, or an alkyl group containing 1-5 carbon atoms, which is optionally substituted with up to 3 =O and/or OH groups, and in which optionally a carbon atom is replaced with an S atom,

Z₃ is, in each case independently, =O, OH, NO₂, NH₂, halogen, or an alkyl group containing 1-5 carbon atoms, which is optionally substituted with up to 3 =O and/or OH groups, and in which optionally a carbon atom is replaced with an O atom, and

n is, in each case independently, 0, 1, 2 or 3;



wherein,

G_1 to G_7 are, in each case independently, C, O, S, or N, wherein at least 3 of G_1 to G_7 are C atoms,


Z_1 is, in each case independently, absent, or $=O$, $=NH$ or an alkyl group containing 1 to 5 carbon atoms,

Z_2 is, in each case independently, a straight chain or branched alkyl group containing 1 to 5 carbon atoms, in which optionally a carbon atom is replaced with an O atom, and which is optionally substituted with 1 or 2 $=O$ and/or $-OH$ groups,

Q is, in each case independently, a bond or an alkylene group containing 1 to 5 carbon atoms, which is optionally substituted with $=O$, in which optionally 1, 2, or 3 carbon atoms are, in each case independently, replaced with an N or S atom, wherein S is optionally substituted with 1 or 2 $=O$ groups, and

n is 0, 1 or 2, and

m is 1 or 2;



wherein,

A is a 5-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2 or 3 heteroatoms selected from O, S, and N, or is a C_{10} aromatic bi-cyclic ring containing 1, 2 or 3 heteroatoms selected from O, S, and N,

~~D is absent or is a fully or partially saturated or unsaturated cyclic ring containing 6 or 7 carbon atoms;~~


~~Q₁ and Q₂ are, each independently of each other, a bond or a straight chain or branched alkylene group containing 1-10 carbon atoms, which is optionally substituted with an =O group, and in which optionally 1, 2 or 3 carbon atoms, independently of each other, are replaced with an N or O atom, and wherein optionally 1-3 carbon atoms are replaced with a C= and/or =C, and/or when the alkylene group is straight chain with a phenyl group, wherein Q₂ is absent when D is absent;~~

~~Z₁ is, in each case independently, absent or an alkyl group containing 1 to 5 carbon atoms, in which optionally a carbon atom is replaced with an O group, and which is optionally substituted with one or two =O or OH groups;~~

~~Z₂ is, in each case independently, =O or an alkyl group containing 1-5 carbon atoms, which is optionally substituted with =O in one or two places and/or OH;~~

~~Z₃ is halogen, or an alkyl group containing 1 to 5 carbon atoms, which is optionally halogenated, and~~

~~n is 1 or 2;~~




~~wherein,~~

~~A is a 5- or 6- membered saturated or partially or fully unsaturated heterocyclic ring containing 2 or 3 heteroatoms selected from S and N;~~

~~Z is, in each case independently, a straight chain or branched alkyl group containing 3-5 carbon atoms, which is substituted with =O and/or OH groups, and in which a carbon atom is replaced with an S atom, and~~

~~n is 1, 2, or 3;~~



wherein,

G_1 to G_{12} are, each independently of each other, C, N, S or O,


Z is, in each case independently, an alkyl containing 1 to 5 carbon atoms, which is optionally substituted with 1 to 2 =O and/or -OH groups,

Q is a bond or an alkylene group containing 1 to 5 carbon atoms,

m is 0, 1, 2 or 3,

n is 0, 1, 2 or 3, such that

$m+n \geq 1$;



wherein,


Z_1 is, in each case independently, halogen, NO_2 or -OH,

Z_2 is, in each case independently, an alkyl group containing 1-5 carbon atoms, which is optionally substituted with an =O and/or -OH group, and in which optionally a carbon atom is replaced with an S atom,

n is 0, 1, 2, or 3,

m is 0, 1, 2, or 3, and


$n+m$ is 3 or more;



wherein,

Z is, in each case independently, $C\equiv N$, halogen, or an alkyl group containing 1-5 carbon atoms, which is optionally substituted with halogen, and/or is substituted with one or more $=O$ and/or $-OH$ groups, and in which optionally a carbon atom is replaced with an S atom, and

n is 2, 3, 4 or 5;



wherein,

D_1 is a 5- or 6- membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2, or 3 heteroatoms selected from O, S, and N,

D_2 is a 6- membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2, or 3 heteroatoms selected from O, S, and N, or is optionally a phenylene group when D_3 is present,

D_3 is absent or a 5- or 6- membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2, or 3 heteroatoms selected from O, S, and N,

Q_1 is $-O-$, or a straight chain alkylene group containing 1-5 carbon atoms, in which optionally a carbon atom is replaced with an N, O or S atom, and which is optionally substituted with an $=O$ atom,


Q_2 is absent when D_3 is absent or is a bond or an $-O-$ group,

Z_1 is, in each case independently, $=O$ or halogen,

Z_2 is, in each case independently, $=O$, $C\equiv N$, $COOH$, NO_2 or halogen,

Z_3 is, in each case independently, halogen, and is absent when D_3 is absent, and

n is, in each case independently, 0, 1, 2, or 3;



wherein,


B is a phenylene group,

Q is a straight chain alkylene group containing 1-10 carbon atoms, in which optionally up to three carbon atoms are replaced with an N, O or S atom, and which is optionally substituted with 1 or 2 $=O$ groups,

Z is, in each case independently, halogen, or an alkyl group containing 1-5 carbon atoms, in which optionally a carbon atom is replaced with an O atom,

n is 0 or 1, and

m is 1 or 2;




wherein,

D is, a 6-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2, or 3 heteroatoms selected from O, S, and N,

Z is $=O$

n is 1, or 2;




wherein,

Q is, each independently, C or N, wherein,

Z_1 is a phenyl group, or 2 of Z_1 together form with the Q atoms to which they are bound a 6-membered aromatic ring containing only C atoms,

Z_2 is halogen, preferably Cl, and

n is 1, or 2;




wherein,

D is, a carbocyclic group containing 8 to 10 carbon atoms, and


R is OH or an alkyl group containing 1 to 5 carbon atoms, in which optionally a carbon atom is replaced with an N or O atom or with a phenyl group, and which is optionally substituted with 1 to 2 =O and/or OH groups.

3. (withdrawn) A pharmaceutical composition comprising a compound of formula I to XVII or a pharmaceutically acceptable salt thereof



wherein,

- B is a phenyl ring,
 D is a phenyl ring or a 5-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2 or 3 heteroatoms selected from O, S, and N,
 A is, in each case independently of each other, a 5-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2 or 3 heteroatoms selected from O, S, and N,
 Q is a bond or an alkylene or alkenylene group containing 1-5 carbon atoms, which is optionally substituted with $=O$, and in which optionally a carbon atom is replaced with an N atom,
 Z_1 is, in each case independently, $-NH_2$, $=O$, $=NH$, or $=N$ -phenyl, -phenyl, or alkyl containing 1 to 5 carbon atoms,
 Z_2 is, in each case independently, $-OH$, halogen, alkyl containing 1-5 carbon atoms, which is optionally substituted with halogen, and/or substituted with $=O$ and/or $-OH$, and in which one C atom is optionally replaced with an O atom,
 Z_3 is, in each case independently, alkyl containing 1-5 carbon atoms, and
 n is, in each case independently, 0, 1, 2, or 3;



wherein,

G_1 , G_2 , and G_3 are, in each case independently, C, O, S, or N,

D is a phenyl ring or a 5- or 6-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2 or 3 heteroatoms selected from O, S, and N,


Q is a straight chain or branched alkylene or alkenylene group containing 1-5 carbon atoms, which is optionally substituted with =O and/or -OH, and in which optionally 1 or 2 carbon atoms, independently of each other, are replaced with an N or S atom,

Z_1 is, in each case independently, =O, halogen, or an alkyl group containing 1-5 carbon atoms, which is optionally substituted with =O and/or -OH,

Z_2 is, in each case independently, =O, or an alkyl group containing 1-5 carbon atoms, which is optionally substituted with =O and/or -OH,

n is 0, 1, or 2, and

m is 0, or 1;




wherein,

G_1 , G_2 , G_3 , G_4 , G_5 and G_6 are, in each case independently, C, O, S, or N, such that four or five of G_1 , G_2 , G_3 , G_4 , G_5 and G_6 are C atoms and the remaining G_1 , G_2 , G_3 , G_4 , G_5 and G_6 are O, S, or N,

Q is a bond or a straight chain or branched alkylene or alkenylene group containing 1-10 carbon atoms which is optionally substituted with =O in one or two places, and in which optionally 1 or 2 carbon atoms, independently of each other, are replaced with an N or O atom, and in which optionally a carbon atom is replaced with a 6-membered heterocyclic group containing 1 or 2 nitrogen atoms when the alkylene or alkenylene group is a straight chain group,

- Z_1 is, in each case independently, -OH, halogen, or an alkyl group containing 1-5 carbon atoms,
- Z_2 is, in each case independently, =O, halogen, or an alkyl group containing 1-10 carbon atoms which is optionally substituted with =O in one or two places and/or -OH, and in which optionally 1 or 2 carbon atoms, independently of each other, are replaced with an N and/or S atom,
- Z_3 is, in each case independently, -OH, halogen, -NO₂, an alkyl group containing 1-5 carbon atoms, in which optionally a carbon atom is replaced with an O atom, and which is optionally substituted with =O in one or two places, or is -O-phenyl, wherein the phenyl group in the -O-phenyl is optionally substituted with an -NO₂ group,
- n is 0, 1, or 2, and
- m is 0, 1, 2, or 3;




(IV)

wherein,

- B is a phenyl ring,
- D₁ is a phenylene ring or a 6-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2, or 3 heteroatoms selected from O, S, and N,
- D₂ and D₂' are, each independently of each other, absent or a phenyl or phenylene ring or a 5- or 6-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2 or 3 heteroatoms selected from O, S, and N,
- Q₁ is a bond or a branched or straight chain alkylene or alkenylene group containing 1-10 carbon atoms, which is optionally substituted with 1 to 5 =O and/or OH groups, in which optionally 1, 2, or 3 carbon atoms are, in each case independently, replaced with an N, O or S atom, wherein S is optionally substituted with 1 or 2 =O groups,
- Q₂ and Q₂' are, each independently of each other, a bond or a branched or straight chain


alkylene group containing 1-5 carbon atoms, which is optionally substituted with an =O group, in which optionally a carbon atom is replaced with an N, S, or O atom, wherein Q_2 is absent when D_2 is absent and Q_2' is absent when D_2' is absent,

- Z is, in each case independently, =O, =S, -OH, -NH₂, -NO₂, -C≡N, -SO₃H, is halogen, or a straight chain or branched alkyl or alkenyl group containing 1 to 10, which is optionally substituted with 1 to 3 =O and/or OH groups, and in which optionally a carbon atom is replaced with an N, O or S atom, or is a cyclic alkyl group containing 3 carbon atoms,
- n is, in each case independently, 0, 1, 2, 3, 4 or 5;



wherein,


- Z is, in each case independently, -NO₂, an alkyl containing 1 to 5 carbon atoms, in which optionally a carbon atom is replaced with an O atom, and which is optionally substituted with an =O group,
- A is a straight chain alkylene group containing 1 to 5 carbon atoms, and
- n is 1, 2 or 3;



wherein,

- B is a phenyl ring,
- D is absent, or is a phenyl ring or a 6-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2, or 3 heteroatoms selected from O, S, and N,

- A is a 5-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms selected from O, S, and N,
- Q₁ and Q₂ are, in each case independently of each other, a bond or a straight chain or branched alkylene group containing 1-5 carbon atoms, in which optionally a carbon atom is replaced with an O, N or S atom, and in which optionally 1 or 2 -C- groups are replaced with -C= or =C- groups, and which is optionally substituted with an =O group, wherein Q₂ is absent when D is absent,
- Z₁ is, in each case independently, -NO₂, -OH, halogen, or an alkyl group containing 1-5 carbon atoms, which is optionally substituted with up to 3 =O and/or OH groups, and in which optionally a carbon atom is replaced with and N, S or O atom,
- Z₂ is, in each case independently, -NH₂, -OH, =NH, =O, =S, phenyl, halogen, or an alkyl group containing 1-5 carbon atoms, which is optionally substituted with up to 3 =O and/or OH groups, and in which optionally a carbon atom is replaced with an S atom,
- Z₃ is, in each case independently, =O, -OH, NO₂, NH₂, halogen, or an alkyl group containing 1-5 carbon atoms, which is optionally substituted with up to 3 =O and/or OH groups, and in which optionally a carbon atom is replaced with an O atom, and
- n is, in each case independently, 0, 1, 2 or 3;



wherein,


- G₁ to G₇ are, in each case independently, C, O, S, or N, wherein at least 3 of G₁ to G₇ are C atoms,
- Z₁ is, in each case independently, absent, or =O, =NH or an alkyl group containing 1 to 5 carbon atoms,

Z_2 is, in each case independently, a straight chain or branched alkyl group containing 1 to 5 carbon atoms, in which optionally a carbon atom is replaced with an O atom, and which is optionally substituted with 1 or 2 =O and/or -OH groups,

Q is, in each case independently, a bond or an alkylene group containing 1-5 carbon atoms, which is optionally substituted with =O, in which optionally 1, 2, or 3 carbon atoms are, in each case independently, replaced with an N or S atom, wherein S is optionally substituted with 1 or 2 =O groups, and

n is 0, 1 or 2, and

m is 1 or 2;



wherein,

A is a 5-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2 or 3 heteroatoms selected from O, S, and N, or is a C_{10} aromatic bi-cyclic ring containing 1, 2 or 3 heteroatoms selected from O, S, and N,

D is absent or is a fully or partially saturated or unsaturated cyclic ring containing 6 or 7 carbon atoms,

Q_1 and Q_2 are, each independently of each other, a bond or a straight chain or branched alkylene group containing 1-10 carbon atoms, which is optionally substituted with an =O group, and in which optionally 1, 2 or 3 carbon atoms, independently of each other, are replaced with an N or O atom, and wherein optionally 1-3 carbon atoms are replaced with a -C= and/or =C-, and/or when the alkylene group is straight chain with a phenyl group, wherein Q_2 is absent when D is absent,


Z_1 is, in each case independently, absent or an alkyl group containing 1 to 5 carbon atoms, in which optionally a carbon atom is replaced with an -O- group, and which is optionally substituted with one or two =O or -OH groups,

- Z_2 is, in each case independently, =O or an alkyl group containing 1-5 carbon atoms, which is optionally substituted with =O in one or two places and/or -OH,
- Z_3 is halogen, or an alkyl group containing 1 to 5 carbon atoms, which is optionally halogenated, and
- n is 1 or 2;




wherein,

- A is a 5- or 6- membered saturated or partially or fully unsaturated heterocyclic ring containing 2 or 3 heteroatoms selected from S and N,
- Z is, in each case independently, a straight chain or branched alkyl group containing 3-5 carbon atoms, which is substituted with =O and/or OH groups, and in which a carbon atom is replaced with an S atom, and
- n is 1, 2, or 3;



wherein,

- G_1 to G_{12} are, each independently of each other, C, N, S or O,
- Z is, in each case independently, an alkyl containing 1 to 5 carbon atoms, which is optionally substituted with 1 to 2 =O and/or -OH groups,
- Q is a bond or an alkylene group containing 1 to 5 carbon atoms,
- m 0, 1, 2 or 3,
- n 0, 1, 2 or 3, such that
- $m+n \geq 1$;



wherein,


Z_1 is, in each case independently, halogen, $-\text{NO}_2$ or $-\text{OH}$,

Z_2 is, in each case independently, an alkyl group containing 1-5 carbon atoms, which is optionally substituted with an $=\text{O}$ and/or $-\text{OH}$ group, and in which optionally a carbon atom is replaced with an S atom,

n is 0, 1, 2, or 3,

m is 0, 1, 2, or 3, and


$n + m$ is 3 or more;



wherein,


Z is, in each case independently, $-\text{C}\equiv\text{N}$, halogen, or an alkyl group containing 1-5 carbon atoms, which is optionally substituted with halogen, and/or is substituted with one or more $=\text{O}$ and/or $-\text{OH}$ groups, and in which optionally a carbon atom is replaced with an S atom, and

n is 2, 3, 4 or 5;




wherein,

- D_1 is a 5- or 6-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2, or 3 heteroatoms selected from O, S, and N,
- D_2 is a 6-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2, or 3 heteroatoms selected from O, S, and N, or is optionally a phenylene group when D_3 is present,
- D_3 is absent or a 5- or 6-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2, or 3 heteroatoms selected from O, S, and N,
- Q_1 is -O-, or a straight chain alkylene group containing 1-5 carbon atoms, in which optionally a carbon atom is replaced with an N, O or S atom, and which is optionally substituted with an =O atom,
- Q_2 is absent when D_3 is absent or is a bond or an -O- group,
- Z_1 is, in each case independently, =O or halogen,
- Z_2 is, in each case independently, =O, $-C\equiv N$, $-COOH$, $-NO_2$ or halogen,
- Z_3 is, in each case independently, halogen, and is absent when D_3 is absent, and
- n is, in each case independently, 0, 1, 2, or 3;




wherein,

- B is a phenylene group,
- Q is a straight chain alkylene group containing 1-10 carbon atoms, in which optionally up to three carbon atoms are replaced with an N, O or S atom, and which is optionally substituted with 1 or 2 =O groups,
- Z is, in each case independently, halogen, or an alkyl group containing 1-5 carbon atoms, in which optionally a carbon atom is replaced with an O atom,
- n is 0 or 1, and
- m is 1 or 2;




wherein,

- D is, a 6-membered saturated or partially or fully unsaturated heterocyclic ring containing 1, 2, or 3 heteroatoms selected from O, S, and N,
- Z is =O
- n is 1, or 2;



wherein,

- Q is, each independently, C or N, wherein,
 Z₁ is a phenyl group, or 2 of Z₁ together form with the Q atoms to which they are bound a 6-membered aromatic ring containing only C atoms,
 Z₂ is halogen, preferably Cl, and
 n is 1, or 2;




















(XVII)


wherein,

- D is, a carbocyclic group containing 8 to 10 carbon atoms, and
 R is -OH or an alkyl group containing 1 to 5 carbon atoms, in which optionally a carbon atom is replaced with an N or O atom or with a phenyl group, and which is optionally substituted with 1 to 2 =O and/or -OH groups.

4. (withdrawn) A method of claim 1, wherein immunosuppression is affected.
5. (withdrawn) A method of claim 1, wherein said patient suffers from an autoimmune disease or from transplant rejection.
6. (withdrawn) A method of claim 5, wherein said patient suffers from rheumatoid arthritis.
7. (original) A method of claim 1, wherein said patient suffers from a neoplasm or a hyperplasia.


8. (original) A method of claim 7, wherein said patient suffers from a benign or malignant tumor.
9. (withdrawn) A method of claim 1, wherein said patient suffers from a depressed immune system.
10. (currently amended) A method of claim 1, wherein said patient suffers from leukemia, lymphoma, ovarian cancer or ~~and~~ breast cancer.
11. (original) A method of claim 1, wherein said patient is human.
12. (currently amended) A method of claim 1, wherein one of the following compounds or a pharmaceutically acceptable salt thereof is administered











OF


















13. (currently amended) A method of claim 2, wherein one of the following compounds or a pharmaceutically acceptable salt thereof is administered







OF

14. (withdrawn) A pharmaceutical composition according to claim 3, comprising one of the following compounds or a pharmaceutically acceptable salt thereof







213




220




222




232




239




245




248




249




254




259




261




262




264




275



276



285




287

or

15. (withdrawn) A method of claim 12, wherein the compound 73 or 92 or a pharmaceutically acceptable salt thereof is administered.
16. (withdrawn) A method of claim 13, wherein the compound 73 or 92 or a pharmaceutically acceptable salt thereof is administered.
17. (withdrawn) A pharmaceutical composition according to claim 14, comprising the compound 73 or 92 or a pharmaceutically acceptable salt thereof.
18. (currently amended) A method of claim 1, wherein the compound of formula I ~~formulae I to XVII~~ has a solubility such that the ClogP value is ≤ 5 , a molecular weight of ≤ 500 Daltons, and ≤ 10 hydrogen bond donors and acceptors.
19. (currently amended) A method of claim 2, wherein the compound of formula I ~~formulae I to XVII~~ has a solubility such that the ClogP value is ≤ 5 , a molecular weight of ≤ 500 Daltons, and ≤ 10 hydrogen bond donors and acceptors.
20. (withdrawn) A pharmaceutical composition according to claim 3, wherein the compound of formulae I to XVII has a solubility such that the ClogP value is ≤ 5 , a molecular weight of ≤ 500 Daltons, and ≤ 10 hydrogen bond donors and acceptors.
21. (currently amended) A method according to claim 1 comprising administering an effective amount of a compound formula I ~~formulae I to IX~~ or a pharmaceutically acceptable salt thereof.
22. (original) A method according to claim 2 comprising administering a compound of formula I ~~formula I to IX~~ or a pharmaceutically acceptable salt thereof.
23. (withdrawn) A pharmaceutical composition according to claim 3 comprising a

compound of formula I to IX or a pharmaceutically acceptable salt thereof.

24. (new) A method of claim 1, wherein the compound is



25. (new) A method of claim 1, wherein the compound is

